


-40-

CLAIMS

1. A prodrug having the formula



5 the stereoisomeric form or salt thereof, wherein

n is 1, 2, 3, 4 or 5;

Y is proline, alanine, hydroxyproline, dihydroxyproline, thiazolidinecarboxylic acid (thiopline), dehydroproline, pipecolic acid (L-homopline),
 10 azetidinecarboxylic acid, aziridinecarboxylic acid, glycine, serine, valine,
 leucine, isoleucine and threonine;

X is selected from any amino acid in the D- or L-configuration;

X and Y in each repeat of [Y-X] are chosen independently from one another and
 independently from other repeats;

15 Z is a direct bond or a bivalent straight or branched saturated hydrocarbon group
 having from 1 to 4 carbon atoms;

R¹ is an aryl, heteroaryl, aryloxy, heteroaryloxy, aryloxyC₁₋₄alkyl,
 heterocycloalkyloxy, heterocycloalkylC₁₋₄alkyloxy, heteroaryloxyC₁₋₄alkyl,
 heteroarylC₁₋₄alkyloxy;

R² is arylC₁₋₄alkyl;

20 R³ is C₁₋₁₀alkyl, C₂₋₆alkenyl or C₃₋₇cycloalkylC₁₋₄alkyl;

R⁴ is hydrogen or C₁₋₄alkyl;

aryl, when used alone or in combination with another group, means phenyl
 optionally substituted with one or more substituents each individually selected
 from the group consisting of C₁₋₄alkyl, hydroxy, C₁₋₄alkyloxy, nitro, cyano,
 25 halo, amino, mono- or di(C₁₋₄alkyl)amino and amido;

heteroaryl, when used alone or in combination with another group, means a
 monocyclic or bicyclic aromatic heterocycle having one or more oxygen,
 sulphur or nitrogen heteroatoms, which aromatic heterocycle may optionally
 be substituted on one or more carbon atoms with a substituent selected from
 the group consisting of C₁₋₄alkyl, C₁₋₄alkyloxy, amino, hydroxy, aryl, amido,
 30 mono- or di(C₁₋₄alkyl)amino, halo, nitro, heterocycloalkyl and C₁₋₄
 alkyloxycarbonyl, and which aromatic heterocycle may also be optionally
 substituted on a secondary nitrogen atom by C₁₋₄alkyl or arylC₁₋₄alkyl;


-41-

- heterocycloalkyl, when used alone or in combination with another group, means a saturated or partially unsaturated monocyclic or bicyclic heterocycle having one or more oxygen, sulphur or nitrogen heteroatoms, which heterocycle may optionally be substituted on one or more carbon atoms with a substituent
5 selected from the group consisting of C₁₋₄alkyl, C₁₋₄alkyloxy, hydroxy, halo and oxo, and which heterocycle may also be optionally substituted on a secondary nitrogen atom by C₁₋₄alkyl or arylC₁₋₄alkyl.
2. A prodrug as claimed in claim 1 wherein each X independently is selected from a naturally occurring amino acid.
- 10 3. A prodrug as claimed in claim 1 or 2 wherein n is 1, 2 or 3.
4. A prodrug as claimed in any one of claims 1 to 3 wherein n is 2 or 3 and wherein at least one X is an hydrophobic or aromatic amino acid.
5. A prodrug as claimed in any one of claims 1 to 4 wherein n is 2 or 3 and wherein at least one X is an neutral or acidic amino acid.
- 15 6. A prodrug as claimed in any one of claims 1 to 5 wherein n is 2 or 3 and wherein at least one X is a basic amino acid.
7. A prodrug as claimed in any one of claims 1 to 6 wherein -(Y-X)_n comprises amino-terminally X-Pro, X-Ala, X-Gly, X-Ser, X-Val, or X-Leu.
8. A prodrug as claimed in any one of claims 1 to 7 wherein -(Y-X)_n comprises
20 amino-terminally X-proline or X-alanine.
9. A prodrug as claimed in any one of claims 1 to 8 wherein each Y independently is proline, alanine, glycine, serine, valine or leucine.
10. A prodrug as claimed in any one of claims 1 to 9 wherein each Y independently is proline or hydroxyproline or dihydroxyproline or alanine.
- 25 11. A prodrug as claimed in any one of claims 1 to 10 wherein each Y independently is proline or alanine.
12. A prodrug as claimed in any one of claims 1 to 11 wherein -(Y-X)_n is -(Y-X)_{1or2}-Y-Val.
13. A prodrug as claimed in any one of claims 1 to 12 wherein -(Y-X)_n is
30 -(Y-X)_{1or2}-Pro-Val.
14. A prodrug as claimed in any one of claims 1 to 13 wherein the (Y-X)_n oligopeptide is built up with (Y-X) repeats selected from the group consisting of Pro-Val,

-42-


Pro-Asp, Pro-Ser, Pro-Lys, Pro-Arg, Pro-His, Pro-Phe, Pro-Ile, Pro-Leu, Ala-Val, Ala-Asp, Ala-Ser, Ala-Lys, Ala-Arg, Ala-His, Ala-Phe, Ala-Ile and Ala-Leu.

15. A prodrug as claimed in any one of claims 1 to 14 wherein R^1 is heterocycloalkyloxy, heteroaryl, heteroaryl C_{1-4} alkyloxy, aryl or aryloxy C_{1-4} alkyl.
- 5 16. A prodrug as claimed in any one of claims 1 to 15 wherein R^1 is hexahydrofuro[2,3-b]furan-3-yl-oxy, tetrahydrofuran-3-yl-oxy, quinolin-2-yl, thiazol-5-ylmethyloxy, 3-hydroxy-2-methyl-1-phenyl, 2,6-dimethylphenoxyethyl.
17. A prodrug as claimed in any one of claims 1 to 16 wherein R^1 is hexahydrofuro[2,3-b]furan-3-yl-oxy, tetrahydrofuran-3-yl-oxy, quinolin-2-yl,
- 10 18. A prodrug as claimed in any one of claims 1 to 17 wherein R^1 is (3R, 3aS, 6aR)-thiazol-5-ylmethyloxy, 3-hydroxy-2-methyl-1-phenyl, 2,6-dimethylphenoxyethyl.
19. A prodrug as claimed in any one of claims 1 to 18 wherein R^2 is phenylmethyl; R^3 is isobutyl and R^4 is hydrogen.
- 15 20. A prodrug as claimed in any one of claims 1 to 19 wherein Z is methylene.
21. A prodrug according to claim 1 wherein the prodrug is




or a salt thereof.

22. A prodrug according to claim 1 wherein the prodrug is



or a salt thereof.


- 20 23. A prodrug according to claim 1 wherein the prodrug is



or a salt thereof.


24. A prodrug according to any one of claims 1 to 23 for use as a medicine.

25. Use of a prodrug according to any one of claims 1 to 23 for the manufacture of a medicament useful for preventing or treating HIV infection.
26. A method of preventing or treating HIV infection by administering to any host, including a human, a non-human animal and mammals, a prodrug according to any one of claims 1 to 23 in an amount effective to prevent or treat the HIV infection.
27. A pharmaceutical preparation which contains an effective dose of at least one of the prodrugs as claimed in any one of claims 1 to 23 in addition to customary pharmaceutically innocuous excipients and auxiliaries.
28. A method for modulating the water solubility, modulating plasma protein binding and/or the bioavailability of a therapeutic compound



by coupling a peptide of formula H-(X-Y)_n to said prodrug wherein n, X, Y, R¹, R², R³, R⁴ and Z are as defined in any one of claims 1 to 23 and wherein the resulting conjugate is cleavable by a dipeptidyl-peptidase.

29. A method according to claim 28 wherein the dipeptidyl-peptidase is CD26.
30. A method of producing a prodrug of a therapeutic compound



wherein the prodrug is cleavable by a dipeptidyl-peptidase, the method comprising the step of linking a therapeutic compound and a peptide of formula H-(X-Y)_n wherein n, X, Y, R¹, R², R³, R⁴ and Z are as defined in any one of claims 1 to 20 and wherein the resulting conjugate is cleavable by a dipeptidyl-peptidase.

31. A method according to claim 30 wherein the dipeptidyl-peptidase is CD26.